arXiv:cond-mat/9907042v1 [cond-mat.stat-mech] 2 Jul 1999

THERMODYNAMIC LIMIT IN CHERN-SIMONS SYSTEM OF PARTICLES WITH MB STATISTICS.

W.I.Skrypnik

Institute of mathematics, Tereshchenkivska str.3, Kyiv-4, Ukraine, 252601

Abstract

The reduced density matrices (RDMs) are calculated in the thermodynamic limit for the Chern-Simons non-relativistic particle system and Maxwell-Boltzmann (MB) statistics. It is established that they are zero outside of a diagonal and well-behaved after a renormalization, depending on an arbitrary real number, if the condition of neutrality holds.

1 Introduction.

The Chern-Simons 2-d quantum system of n nonrelativistic spinless identical particles of unit mass is described by the Hamiltonian $\dot{H}_n$, defined on $C^\infty(\mathbb{R}^{2n}\setminus \cup_{j,k}(x_j = x_k))$

$$\dot{H}_n = \frac{1}{2}\sum_{j=1}^{n} ||p_j - a_j(X_n||^2, \qquad a_j^\nu(X_n) = \epsilon^{\nu\mu}\partial_{\mu,j}U_C(X_n), \quad \nu,\mu = 1,2, \tag{1.1}$$

$$U_C(X_n) = g\sum_{1\le k<j\le n} e_je_k ln|x_j - x_k|, \quad X_n = (x_1,...x_n)\in\mathbb{R}^{2n}, \quad p_j^\mu = i^{-1}\partial_{\mu,j} = i^{-1}\frac{\partial}{\partial x_j^\mu},$$

where $||v||^2 = (v^1)^2 + (v^2)^2$, ϵ is the skew symmetric tensor and the repeating index implies a summation, real number e_j(a charge) takes values in a finite set $E_{\{c\}}$ from $\mathbb{R}$.

Differentiating the equality $f(f^{-1}(x)) = x$ we derive the formula

$$\frac{df^{-1}(x)}{dx} = (\frac{df(y)}{dy})^{-1}, \qquad y = f^{-1}(x).$$

From this equality for $f(x) = \tan x$ and the equality $\frac{d}{dx}\tan x = 1 + \tan^2(x)$ the following relation is derived

$$\frac{\partial}{\partial x^\nu}\arctan\frac{x^2}{x^1} = \epsilon^{\nu\mu}x^\mu|x|^{-2} = \epsilon^{\nu\mu}\frac{\partial}{\partial x^\mu}ln|x|.$$

This means that CS system is almost(quasi-)integrable, that is

$$\dot{H}_n = e^{i\hat{U}}\dot{H}_n^0 e^{-i\hat{U}},$$

where $-2\dot{H}_n^0$ is the 2n-dimensional Laplacian $-2H_n^0$ restricted to $C^\infty(\mathbb{R}^{2n}\setminus\cup_{j,k}(x_j = x_k))$ and $\hat{U}$ is the operator of multiplication by $U(X_n)$,

$$U(X_n) = g\sum_{1\le k<j\le n} e_je_k\phi(x_j - x_k), \qquad \phi(x_j - x_k) = \arctan\frac{x_j^2 - x_k^2}{x_j^1 - x_k^1}$$

As a result, there exists the simplest sefadjoint extension H_n of $\dot{H}_n$

with the domain $D(H_n) = e^{i\hat{U}} D(H_n^0)$. Another selfadjoint extension of $\dot{H}_n$ is given by (1.2) in which, instead of H_n^0, another selfadjoint extension of $\dot{H}_n^0$ is considered.

The CS system of particles with different statistics has been studied by many authors [1-4] since it can be derived (formally) in 3-d topological electrodynamics (its Lagrangian contains CS term) in the limit of the vanishing Maxwell term (the same is true for its relativistic version). There is a hope it can give a new mechanism of superconductivity, superfluidity and P, T violation (general outlook of modern statistical mechanics allows to expect breakdown only of discrete (hidden) symmetries for quantum systems at nonzero temperature in 3 space-time dimensions).

In this paper the CS system is considered with the Hamiltonian (1.2) and the Maxwell-Boltzmann (MB) statistics. Its thermodynamics coincides with thermodynamics of the free particle system if Dirichlet boundary condition is considered(we do this in this paper). Our calculations show that the Gibbs (grand-canonical)reduced density matrices (RDMs) $\rho^{B_L}(X_m|Y_m)$ in the sphere B_L of radius L, centered at the origin, tend to zero if $x_j \neq y_j$. They are expressed as the product of a function having a non-zero limit and

$$\exp\{-\frac{N_{r,L}}{4} g^2 (\sum_e z_e e^2) \sum_{j=1}^m e_j^2 |x_j - y_j|^2\}, \qquad N_{r,L} = \int\limits_{r<|x|\leq L} P^{\beta}_{0(B_L)}(x|x)|x|^{-2} dx.$$

where z_e is the activity of the particles with the charge e and $N_{r,L}$ diverges as $2lnL$, $P^{\beta}_{0(B_L)}(x|y)$ is the kernel of the semigroup whose infinitesimal generator coincides with the two-dimensional Laplacian with the Dirichlet boundary condition on the boundary of the ball B_L. We propose to renormalize the RDMs by dividing them by this exponent. We show then the important feature of this renormalization: if the condition of the neutrality holds then the resulting RDMs in the thermodynamic limit $\rho(X_m|Y_m)$ satisfy the compatibility condition(see the corollary), that is, they define a state of the infinite particle system.

The unusual property of the RDMs to be zero almost everywhere was found by us in 1-d (integrable) systems of particles with the Hamiltinian (1.1) when a_j is expressed through a pair (scalar magnetic) long-range potential [5-6]. We expect that the similar property is true for bosonic and fermiomic CS systems for small values of activities of particles(see also [7]).

The results of our calculations are formulated in the theorem and corollary in the next paragragh.

2 Main result.

Let $\Lambda \in \mathbb{R}^2$ be a compact set and assume the Dirichlet boundary conditions on the boundary $\partial\Lambda$. For the inverse tempetature β, and the activity z_{e_s} of the particles with the charge e_s, the Gibbs (grand-canonical, equilibrium) RDMs are given by

$$\rho^{\Lambda}(X_m|Y_m) =$$

$$= Z_{(e)_m} \Xi_{\Lambda}^{-1} \sum_{n\geq 0} \prod_{s=1}^{n} \sum_{e'_s} \frac{z_{e'_s}^{n_s}}{n_s!} \int\limits_{\Lambda^n} dX'_n \exp\{i[U(X_m, X'_n) - U(Y_m, X'_n)]\} P^{\beta}_{0(\Lambda)}(X_m, X'_n|Y_m, X'_n),$$

where Ξ_{Λ} is the grand partition function (it coincides with the numerator in the r.h.s. of this equality for the case "m=0", i.e. when there are no X_m and Y_m), $P^{\beta}_{0(\Lambda)}(X_m|Y_m)$ is the kernel of the semigroup, whose infinitesimal generator concides with $H^0_{n,\Lambda}$ ($-2H^0_{n,\Lambda}$ is the Laplacian in Λ^n with the Dirichlet boundary condition on the boundary $\partial\Lambda^n$), $Z_{(e)_m} = \prod^{m} z_{e_j}$ and the summation in e_s is performed over $E_{\{c\}}$.

$$P^{\beta}_{0(\Lambda)}(X_n|Y_n) = \prod_{j=1}^{n} P^{\beta}_{0(\Lambda)}(x_j|y_j), \quad X_m = (X^1_m, X^2_m), Y_m = (Y^1_m, Y^2_m) \in \mathbb{R}^{2m}, \tag{2.1}$$

$P^{\beta}_{0(\Lambda)}(x|y)$ is the transition probability of the 2-dimensional free diffusion proccess with the Dirichlet boundary condition on $\partial\Lambda$.

$$P^{\beta}_{0(\Lambda)}(x|y) = \int P^{\beta}_{x,y}(d\omega)\chi_{\Lambda}(\omega),$$

$P_{x,y}(d\omega)$ is the conditional Wiener measure and $\chi_{\Lambda}(\omega)$ is the characteristic function of the paths that are inside Λ.

From the equality

$$U(X_m, X'_n) = U(X_m) + \sum_{j=1}^{m}\sum_{k=1}^{n} \phi(x_j - x'_k)e_j e'_k + U(X'_n)$$

we obtain

$$\rho^{\Lambda}(X_m|Y_m) = Z_{(e)_m}\Xi^{-1}_{\Lambda} \exp\{i[U(X_m) - U(Y_m)]\} P^{\beta}_{0(\Lambda)}(X_m|Y_m) \sum_{n\geq 0}\sum_{e'_s} \frac{z^{n_s}_{e'_s}}{n_s!} \times$$
$$\times \int_{\Lambda^n} dX'_n \prod_{k=1}^{n} \exp\{i\sum_{j=1}^{m} e_j e_k(\phi(x_j - x'_k) - \phi(y_j - x'_k))\} P^{\beta}_{0(\Lambda)}(x'_k|x'_k),$$

As a result

$$\rho^{\Lambda}(X_m|Y_m) = Z_{(e)_m} \exp\{i[U(X_m) - U(Y_m)] + G_{\Lambda}(X_m|Y_m)\} P^{\beta}_{0(\Lambda)}(X_m|Y_m), \tag{2.2}$$

$$G_{\Lambda}(X_m|Y_m) = \sum_{e} z_e \int_{\Lambda} P^{\beta}_{0(\Lambda)}(x|x)[\exp\{i\sum_{j=1}^{m} ee_j(\phi(x_j - x) - \phi(y_j - x))\} - 1]dx. \tag{2.3}$$

Here we used the equality $\Xi_{\Lambda} = \exp\{\sum_{e} z_e \int_{\Lambda} P^{\beta}_{0(\Lambda)}(x|x)dx\}$.

With the help of the equality

$$exp\{i \arctan \frac{x^2}{x^1}\} = \frac{x}{|x|} = (\frac{x}{x^*})^{\frac{1}{2}}, \qquad x = x^1 + ix^2,$$

we derive

$$\exp\{i\sum_{j=1}^{m} ee_j(\phi(x_j - x) - \phi(y_j - x))\} = \prod_{j=1}^{m} \left(\frac{(x - x_j)(x^* - y^*_j)}{(x^* - x^*_j)(x - y_j)}\right)^{\frac{1}{2}gee_j} = G_x(X_m|Y_m).$$

We have to use the Taylor expansions for $|x| < 1$

$$(1 - x)^g = 1 - gx + \frac{g(g-1)}{2}x^2 + \sum_{n\geq 3} C^g_n x^n = \sum_{n\geq 0} C^g_n x^n,$$

$$\left(\frac{x-x_j}{x^*-x_j^*}\right)^{g'} = \left(\frac{x}{x^*}\right)^{g'}\left(\frac{1-\frac{x_j}{x}}{1-\frac{x_j^*}{x^*}}\right)^{g'} = \left(\frac{x}{x^*}\right)^{g'}\{1+g'(-\frac{x_j}{x}+\frac{x_j^*}{x^*})+$$

$$+\frac{g'(g'-1)}{2}(\frac{x_j^2}{x^2}+\frac{x_j^{*2}}{x^{*2}})-g'^2|\frac{x_j}{x}|^2+\sum_{n_1^+n_1^-\geq 3} C^{-g'}_{n_1^+}C^{g'}_{n_1^-}(\frac{x_j^*}{x^*})^{n_1^+}(\frac{x_j}{x})^{n_1^-}\},$$

Applying this formula we deduce

$$\left(\frac{(x-x_j)(x^*-y_j^*)}{(x^*-x_j^*)(x-y_j)}\right)^{g'} =$$

$$= 1 + G'_x(x_j|y_j) + \sum_{n_1^+ +...+n_2^-\geq 3} C^{-g'}_{n_1^+}C^{g'}_{n_1^-}C^{-g'}_{n_2^+}C^{g'}_{n_2^-}(\frac{x_j^*}{x^*})^{n_1^+}(\frac{x_j}{x})^{n_1^-}(\frac{y_j^*}{x^*})^{n_2^-}(\frac{y_j}{x})^{n_2^+}.$$

where

$$G'_x(x_j|y_j) = g'(-\frac{x_j-y_j}{x}+\frac{x_j^*-y_j^*}{x^*})-g'^2(\frac{x_j}{x}-\frac{x_j^*}{x^*})(\frac{y_j}{x}-\frac{y_j^*}{x^*})+$$

$$+\frac{g'(g'-1)}{2}(\frac{x_j^2+y_j^2}{x^2}+\frac{x_j^{*2}+y_j^{*2}}{x^{*2}})-g'^2[|\frac{x_j}{x}|^2+|\frac{y_j}{x}|^2].$$

As a result

$$G_x(X_m|Y_m) = 1 + G^0_{x,e}(X_m|Y_m) + \sum_{j=1}^m G'_x(x_j|y_j), \tag{2.4}$$

where

$$G^0_{x,e}(X_m|Y_m) = \sum_{n_{1,1}^+ +...+n_{2,m}^-\geq 3}\prod_{j=1}^m C^{-g'}_{n_{1,j}^+}C^{g'}_{n_{1,j}^-}C^{-g'}_{n_{2,j}^+}C^{g'}_{n_{2,j}^-}(\frac{x_j^*}{x^*})^{n_{1,j}^+}(\frac{x_j}{x})^{n_{1,j}^-}(\frac{y_j^*}{x^*})^{n_{2,j}^-}(\frac{y_j}{x})^{n_{2,j}^+}.$$

It can be checked that

$$-g'^2(\frac{x_j}{x}-\frac{x_j^*}{x^*})(\frac{y_j}{x}-\frac{y_j^*}{x^*})+\frac{g'^2}{2}(\frac{x_j^2+y_j^2}{x^2}+\frac{x_j^{*2}+y_j^{*2}}{x^{*2}})-g'^2[|\frac{x_j}{x}|^2+|\frac{y_j}{x}|^2] =$$

$$= -g'^2|\frac{x_j-y_j}{x}|^2+\frac{g'^2}{2}(\frac{(x_j-y_j)^2}{x^2}+\frac{(x_j^*-y_j^*)^2}{x^{*2}}). \tag{2.5}$$

This yields

$$G'_x(x_j|y_j) = -g'^2|\frac{x_j-y_j}{x}|^2 + G^-_x(x_j|y_j), \qquad G^-_x(x_j|y_j) =$$

$$= g'(-\frac{x_j-y_j}{x}+\frac{x_j^*-y_j^*}{x^*})+\frac{g'^2}{2}(\frac{(x_j-y_j)^2}{x^2}+\frac{(x_j^*-y_j^*)^2}{x^{*2}})-\frac{g'}{2}(\frac{x_j^2+y_j^2}{x^2}+\frac{x_j^{*2}+y_j^{*2}}{x^{*2}}). \tag{2.6}$$

Let $l_m^+ = max(|x_j|,|y_j|, j=1,...,m)$. Let $\Lambda = B_L$ then

$$G_\Lambda(X_m|Y_m) = \sum z_e\{\int P^{\beta}_{0(\Lambda)}(x|x)[\prod_{j=1}^m\left(\frac{x^*-x_j^*}{}\frac{x-y_j}{}\right)^{\frac{1}{2}gee_j}-1]dx+$$

$$+\int\limits_{2l_m^+\le|x|\le L} P^{\beta}_{0(\Lambda)}(x|x)G^0_x(X_m|Y_m)dx+\int\limits_{2l_m^+\le|x|\le L} P^{\beta}_{0(\Lambda)}(x|x)G'_x(X_m|Y_m)dx\}. \qquad (2.7)$$

For $|x|\ge 2l_m^+$ we have the bound

$$G^0_{x,e}(X_m|Y_m)\le\frac{(2l_m^+)^3}{|x|^3}2^{4|g‘|m}.$$

Here we used the inequalities

$$|(\frac{x_j^*}{x^*})^{n^+_{1,j}}(\frac{x_j}{x})^{n^-_{1,j}}(\frac{y_j^*}{x^*})^{n^-_{2,j}}(\frac{y_j}{x})^{n^+_{2,j}}|\le\frac{(2l_m^+)^3}{|x|^3}2^{-(n^+_{1,j}+n^-_{1,j}+n^+_{2,j}+n^-_{2,j}+)},\quad |C_n^{+(-)g}|\le C_n^{-|g|}>0.$$

After applying them we enlarge the sum in (2.5) to the sum over $(Z^+)^{4m}$.

Since $P_\Lambda(x|x)$ tends to $(2\pi\beta)^{-1}$ the first and the second terms in (2.7) have limits when L tends to infinity. We have only to calculate the third term. Let's show that

$$\int\limits_{r\le|x|\le L} G^-_x(x'|y')dx=0. \qquad (2.8)$$

For arbitrary $r, L, v=v^1+iv^2, B=B_L\backslash B_r$ we have

$$\int\limits_B(\frac{v}{x}-\frac{v^*}{x^*})dx=-2i[v^1\int\limits_B\frac{x^2}{|x|^2}dx-v^2\int\limits_B\frac{x^1}{|x|^2}dx]=0,$$

$$\int\limits_B(\frac{v}{x^2}+\frac{v^*}{x^{*2}})dx=2\{v^1\int\limits_B\frac{(x^1)^2-(x^2)^2}{|x|^4}dx+2v^2\int\limits_B\frac{x^1x^2}{|x|^4}dx\}=0.$$

All the above integrals are zero since the all the functions change signs when either a sign of one of the variables is changed , or a permutation is done.

As a result

$$\int\limits_{r<|x|\le L} G'_x(X_m|Y_m)dx=-\frac{1}{4}g^2(\sum_e z_e e^2)N_{r,L}\sum_{j=1}^m e_j^2|x_j-y_j|^2. \qquad (2.9)$$

The integral in the right-hand-side of this equality diverges as $2lnL$. For $g'=k, k\in\mathbb{Z}$ we obtain (2.4) in which we have to put $C_n^k=0$ for $n>k$, $C_n^k=\frac{k!}{(n-k)!n!}$.

From (2.7), (2.9) we derive

$$G_\Lambda(X_m|Y_m)=\sum_e z_e\{\int\limits_{|x|\le 2l_m^+} P^{\beta}_{0(\Lambda)}(x|x)[\prod_{j=1}^m\left(\frac{x^*-x_j^*}{x-x_j}\frac{x-y_j}{x^*-y_j^*}\right)^{\frac{1}{2}gee_j}-1]dx+$$

$$+\int\limits_{2l_m^+\le|x|\le L} P^{\beta}_{0(\Lambda)}(x|x)G^0_x(X_m|Y_m)dx-\frac{1}{4}g^2(\sum_e z_e e^2)N_{2l_m^+,L}\sum_{j=1}^m e_j^2|x_j-y_j|^2. \qquad (2.10)$$

This equality together with (2.8-9), in which $2l_m^+$ is substituted instead of L, yields for arbitrary $0<r<\infty$

$$G_\Lambda(X_m|Y_m) = \int\limits_{|x|\le r} \sum_e z_e[\prod_{j=1}^m \left(\frac{x^*-x_j^*}{x-x_j}\frac{x-y_j}{x^*-y_j^*}\right)^{\frac{1}{2}gee_j} - 1]P^\beta_{0(B_L)}(x|x)dx+$$

$$+\int\limits_{r<|x|\le L} \sum_e z_e[\prod_{j=1}^m \left(\frac{x^*-x_j^*}{x-x_j}\frac{x-y_j}{x^*-y_j^*}\right)^{\frac{1}{2}gee_j} - 1 - \sum_{j=1}^m G'_x(x_j|y_j)]P^\beta_{0(B_L)}(x|x)dx-$$

$$-\frac{1}{4}g^2(\sum_e z_e e^2)N_{r,L}\sum_{j=1}^m e_j^2|x_j-y_j|^2. \qquad (2.11)$$

Using the equality

$$\lim_{L\to\infty} P^\beta_{0(B_L)}(x|x) = (2\pi\beta)^{-1}$$

we deduce that the function G is well defined

$$G(X_m|Y_m) = \lim_{L\to\infty}[G_\Lambda(X_m|Y_m) + \frac{1}{4}g^2(\sum_e z_e e^2)N_{r,L}\sum_{j=1}^m e_j^2|x_j-y_j|^2] =$$

$$= (2\pi\beta)^{-1}\int\limits_{|x|\le r} \sum_e z_e[\prod_{j=1}^m \left(\frac{x^*-x_j^*}{x-x_j}\frac{x-y_j}{x^*-y_j^*}\right)^{\frac{1}{2}gee_j} - 1]dx+$$

$$+ (2\pi\beta)^{-1}\int\limits_{|x|>r} \sum_e z_e[\prod_{j=1}^m \left(\frac{x^*-x_j^*}{x-x_j}\frac{x-y_j}{x^*-y_j^*}\right)^{\frac{1}{2}gee_j} - 1 - \sum_{j=1}^m G'_x(x_j|y_j)]dx. \qquad (2.12)$$

THEOREM

Let Λ be the ball B_L, centered at the origin, of radius L. Then for all the values of the activities z_e and charges e_j the thermodynamic limit for the RDMs exists and is zero if $x_j \ne y_j$. Moreover, the following equality is valid

$$\rho(X_m|Y_m) = \lim_{L\to\infty}\exp\{g^2\frac{N_{r,L}}{4}\sum_{j=1}^m e_j^2|x_j-y_j|^2\}\rho^{B_L}(X_m|Y_m) =$$

$$= Z_{(e)_m}\exp\{i[U(X_m)-U(Y_m)]\}P_0^\beta(X_m|Y_m)e^{G(X_m|Y_m)},$$

where G is defined by (2.12) and

$$P_0^\beta(X_m|Y_m) = (2\pi\beta)^{-m}\prod_{j=1}^m \exp\{-\frac{|x_j-y_j|^2}{2\beta}\}.$$

Now, let the condition of neutrality holds $\sum\limits_e z_e e = 0$, then from (2.12) it follows that

$$G(X_m|Y_m) = (2\pi\beta)^{-1}\int\limits_{|x|\le r} \sum_e z_e[\prod_{j=1}^m \left(\frac{x^*-x_j^*}{x-x_j}\frac{x-y_j}{x^*-y_j^*}\right)^{\frac{1}{2}gee_j} - 1]+$$

$$+(2\pi\beta)^{-1}\int\{\sum z_e[\prod^m \left(\frac{x^*-x_j^*}{}\frac{x-y_j}{}\right)^{\frac{1}{2}gee_j} - 1]+$$

$$+ g^2(\sum_e z_e e^2)\sum_{j=1}^m e_j^2[|\frac{x_j-y_j}{x}|^2 - \frac{1}{2}(\frac{(x_j-y_j)^2}{x^2} + \frac{(x_j^*-y_j^*)^2}{x^{*2}})]\}dx. \qquad (2.13)$$

From (2.13) it follows that $G(x, X_m|x, Y_m) = G(X_m|Y_m)$. Repeating the above arguments the equality

$$\lim_{L\to\infty}(\pi L^2)^{-1}\int_{B_L}\exp\{i\sum_{j=1}^m ee_j[\phi(x-x_j)-\phi(x-y_j)]\}dx = 1$$

is proved. So, we see that the following corollary is true.

COROLLARY. If the condition of neutrality holds then the compatibility condition for the renormalized RDMs in the thermodynamic limit holds

$$\lim_{L\to\infty}\frac{2\beta}{z_e L^2}\int_{B_L}\rho(x, X_m|x, Y_m)dx = \rho(X_m|Y_m).$$

AKNOWLEDGMENTS.

The research described in this publication was possible in part by the Award number UP1-309 of the U.S. Civilian Research and Development Foundation (CRDF) for independent states of the former Soviet Union.

REFERENCES

1 R.Jackiw,S-Y.Pi, Phys.Rev.D15,42,p.3550, 1990.

2 C.Trugenberger, The anyon fluid in the Bogoliubov approximation, Preprint LA-UR 91-2421, Los Alamos, national laboratory.

3 Y.Kitazawa, H.Murayama, Nucl.Phys.,B338, p. 777, 1990.

4 J.Lykken, J.Sonnenschein, N.Weiss, Theory of anyonic superconductivity. A review. Preprint TAUP-1858-91, Int. Journ. of Mod.Phys., 1991.

5 W.Skrypnik, Ukrainian Math.Journ.,47, 12,p.1686, 1995.

6 W.Skrypnik, Mat.Fiz.,Analiz.,Geom., 4, N1/2, p.248, 1997.

7 W.Skrypnik, Ukrainian Math.Journ.,49, 5,p.691, 1997.